



# TEKNOLOGI INOVATIF PERTANIAN



BADAN PENELITIAN DAN PENGEMBANGAN PERTANIAN  
KEMENTERIAN PERTANIAN  
[www.litbang.pertanian.go.id](http://www.litbang.pertanian.go.id)





## Nilam Varietas Patchoulina 1 Patchoulina 1 Nilam Variety

Inventor : Endang Hadipoentyanti, Amalia, Nursalam Sirait  
dan Sri Yuni Hartati

Balai Penelitian Tanaman Rempah dan Obat

*Indonesian Spice and Medicinal Crops Research Institute*

Status Perlindungan HKI : Pendaftaran Varietas

No: 00404/PPVT/S/2017

IPR Protection Status : Variety registration

No: 00404/PPVT/S/2017

Nilam varietas Patchoulina1 merupakan hasil induksi variasi somaklonal yang memiliki sifat tahan terhadap penyakit layu bakteri (*R. solanacearum*). Varietas ini memiliki potensi hasil terna segar  $2,3 \pm 0,125$  kg/tanaman dan bobot kering  $0,761 \pm 0,036$  kg/tanaman/tahun atau setara dengan produksi terna segar  $36,52 \pm 10,13$  t/ha/tahun, produksi terna kering  $12,67 \pm 3,34$  t/ha/tahun. Produksi minyak  $356,37 \pm 13,76$  kg/ha/tahun, dengan kadar minyak  $2,85 \pm 0,57\%$ , kadar Patchouli Alkohol (PA)  $32,53 \pm 3,81\%$ . Varietas Patchoulina1 dapat dibedakan dari Patchoulina2 karena memiliki bentuk pangkal daun tumpul (obtusus), ujung daun runcing-tumpul (acutus-obtusus), tepi daun bergerigi tumpul (crenatus) dan permukaan daun agak cekung.

The Patchoulina 1 variety was developed through an induction of soma clonal variation having a resistance trait to the bacterial wilt of *R. solanacearum*. Yield potential of this variety is  $2.3 \pm 0.125$  kg of fresh terna/plant with a dry weight of  $0.761 \pm 0.036$  kg/plant/year. It is equivalent to  $36.52 \pm 10.13$  ton fresh terna/ha/year or  $12.67 \pm 3.34$  t dry terna/ha/year. The production of oil is  $356.37 \pm 13.76$  kg/ha year with oil content of  $2.85 \pm 0.57\%$  and Patchouli Alcohol (PA) content of  $32.53 \pm 3.81\%$ . Patchoulina 1 variety can be distinguished from Patchoulina 2 variety from the shape of the leaf whereas is round shape in the base (obtuse), pointed-round at the tip of the leaf (acutus-obtusus), and is serrated in the margin of the leaf (crenatus).